

10 年 Framingham 心血管危险评分

(1) 简介

1994 年，弗明汉心脏研究团队首次提出弗明汉危险评分（Framingham risk score, FRS）公式对心血管危险因素进行评估，为临床医生制定治疗策略提供重要的理论依据。FRS 是一种被国际广泛认可的预测未来 10 年冠心病和其他心血管疾病发生风险的公式。既往有大量研究探讨 FRS 在其他疾病中的研究价值，如卒中，是目前临床上最常用的预测心脑血管事件的工具。

(2) 危险因素

FRS 危险因素包括年龄、收缩压、总胆固醇、高密度脂蛋白胆固醇、吸烟史和糖尿病史。根据 Cox 模型和评分表进行危险评分，其计算公式为：

$$\hat{p} = 1 - S_0(t)^{\exp(\sum_{i=1}^p \beta_i X_i - \sum_{i=1}^p \beta_i \bar{X}_i)}$$

(3) 评分规则

如下图所示，根据评估个体的性别选择不同的评分依据，然后根据个体的年龄、总胆固醇水平（或低密度脂蛋白水平）、高密度脂蛋白水平、收缩压和舒张压水平、是否患有糖尿病与吸烟分别得到 6 个分数，分数相加即可估计出 10 年内患冠心病的平均风险。

(4) 风险分层

对受试者进行 10 年 FRS 风险评估之后，需要对受试者进行风险分层。如果 10 年 FRS 风险 $\geq 20.0\%$ ，视为心血管病高危人群，10 年 FRS 风险为 $10.0\% \sim 20\%$ 视为中危人群，10 年 FRS 风险 $< 10.0\%$ 为低危人群。

(5) 评价网站：

https://m.medsci.cn/scale/show.do?s_id=2&id=395f300e

男性:

Step 1

Years	Age	LDL Pts	Chol Pts
30-34	-9	[-9]	
35-39	-4	[-4]	
40-44	0	[0]	
45-49	3	[3]	
50-54	6	[6]	
55-59	7	[7]	
60-64	8	[8]	
65-69	8	[8]	
70-74	8	[8]	

Step 2

LDL - C			
(mg/dl)	(mmol/L)	LDL Pts	Chol Pts
<100	<2.59	-2	
100-129	2.60-3.36	0	
130-159	3.37-4.14	0	
160-190	4.15-4.92	2	
>190	>4.92	2	

Cholesterol			
(mg/dl)	(mmol/L)	LDL Pts	Chol Pts
<160	<4.14	[-2]	
160-199	4.15-5.17	[0]	
200-239	5.18-6.21	[1]	
240-279	6.22-7.24	[1]	
≥280	≥7.25	[2]	

Step 3

HDL - C			
(mg/dl)	(mmol/L)	LDL Pts	Chol Pts
<40	<1.04	-3	
35-44	0.91-1.16	2	[2]
45-49	1.17-1.29	1	[1]
50-59	1.30-1.55	0	[0]
≥60	≥1.56	-2	[-2]

Step 4

Blood Pressure			
Systolic (mm Hg)	Diastolic (mm Hg)	pts	pts
<120	<80	-3	[-3]
120-129	80-84	0	[0]
130-139	85-89	0	[0]
140-159	90-99	2	[2]
≥160	≥100	3	[3]

Note: When systolic and diastolic pressures provide different estimates for point scores, use the higher number

Step 5

Diabetes			
Diabetes	LDL Pts	Chol Pts	pts
No	0	[0]	
Yes	4	[4]	

Step 6

Smoker			
Smoker	LDL Pts	Chol Pts	pts
No	0	[0]	
Yes	2	[2]	

(sum from steps 1-6)

Step 7

Adding up the points	
Age	_____
LDL-C or Chol	_____
HDL - C	_____
Blood Pressure	_____
Diabetes	_____
Smoker	_____
Point total	_____

(determine CHD risk from point total)

Step 8

CHD Risk			
LDL Pts	10 Yr CHD Risk	Chol Pts	10 Yr CHD Risk
≤-2	1%	[-2]	[1%]
-1	2%	[-1]	[2%]
0	2%	[0]	[2%]
1	2%	[1]	[2%]
2	3%	[2]	[3%]
3	3%	[3]	[3%]
4	4%	[4]	[4%]
5	5%	[5]	[5%]
6	6%	[6]	[6%]
7	7%	[7]	[7%]
8	8%	[8]	[8%]
9	9%	[9]	[9%]
10	11%	[10]	[10%]
11	13%	[11]	[13%]
12	15%	[12]	[15%]
13	17%	[13]	[18%]
14	20%	[14]	[18%]
15	24%	[15]	[20%]
16	27%	[16]	[24%]
≥17	≥30%	≥17	≥27%

(compare to average person your age)

Step 9

Comparative Risk			
Age (years)	Average 10 Yr CHD Risk	Average 10 Yr Hard* CHD Risk	Low** 10 Yr CHD Risk
30-34	<1%	<1%	<1%
35-39	<1%	<1%	1%
40-44	2%	1%	2%
45-49	3%	2%	3%
50-54	8%	3%	5%
55-59	12%	7%	7%
60-64	12%	8%	8%
65-69	13%	8%	8%
70-74	14%	11%	8%

* Hard CHD events exclude angina pectoris

** Low risk was calculated for a person the same age, optimal blood pressure, LDL-C 100-129 mg/dL or cholesterol 160-199 mg/dL, HDL-C ≥45 mg/dL for men or 55 mg/dL for women, non-smoker, no diabetes

Risk estimates were derived from the experience of the Framingham Heart Study, a predominantly Caucasian population in Massachusetts, USA

女性:

Step 1

Years	Age	LDL Pts	Chol Pts
30-34	-1	[-1]	
35-39	0	[0]	
40-44	1	[1]	
45-49	2	[2]	
50-54	3	[3]	
55-59	4	[4]	
60-64	5	[5]	
65-69	6	[6]	
70-74	7	[7]	

Step 2

LDL - C			
(mg/dl)	(mmol/L)	LDL Pts	Chol Pts
<100	<2.59	-3	
100-129	2.60-3.36	0	
130-159	3.37-4.14	0	
160-190	4.15-4.92	1	
>190	>4.92	2	

Cholesterol			
(mg/dl)	(mmol/L)	LDL Pts	Chol Pts
<160	<4.14	[-3]	
160-199	4.15-5.17	[0]	
200-239	5.18-6.21	[1]	
240-279	6.22-7.24	[2]	
≥280	≥7.25	[3]	

Step 3

HDL - C			
(mg/dl)	(mmol/L)	LDL Pts	Chol Pts
<40	<1.04	-3	
35-44	0.91-1.16	1	[1]
45-49	1.17-1.29	0	[0]
50-59	1.30-1.55	0	[0]
≥60	≥1.56	-1	[-1]

Step 4

Blood Pressure			
Systolic (mm Hg)	Diastolic (mm Hg)	pts	pts
<120	<80	0	[0]
120-129	80-84	0	[0]
130-139	85-89	1	[1]
140-159	90-99	2	[2]
≥160	≥100	3	[3]

Note: When systolic and diastolic pressures provide different estimates for point scores, use the higher number

Step 5

Diabetes			
Diabetes	LDL Pts	Chol Pts	pts
No	0	[0]	
Yes	2	[2]	

Step 6

Smoker			
Smoker	LDL Pts	Chol Pts	pts
No	0	[0]	
Yes	2	[2]	

(sum from steps 1-6)

Step 7

Adding up the points	
Age	_____
LDL-C or Chol	_____
HDL - C	_____
Blood Pressure	_____
Diabetes	_____
Smoker	_____
Point total	_____

(determine CHD risk from point total)

Step 8

CHD Risk			
LDL Pts	10 Yr CHD Risk	Chol Pts	10 Yr CHD Risk
<-3	1%		
-2	2%		
-1	2%	[-1]	[2%]
0	3%	[0]	[3%]
1	4%	[1]	[3%]
2	4%	[2]	[4%]
3	6%	[3]	[5%]
4	7%	[4]	[7%]
5	9%	[5]	[8%]
6	11%	[6]	[10%]
7	14%	[7]	[13%]
8	18%	[8]	[16%]
9	22%	[9]	[20%]
10	27%	[10]	[25%]
11	33%	[11]	[31%]
12	40%	[12]	[37%]
13	47%	[13]	[45%]
≥14	≥56%	≥14	≥53%

(compare to average person your age)

Step 9

Comparative Risk			
Age (years)	Average 10 Yr CHD Risk	Average 10 Yr Hard* CHD Risk	Low** 10 Yr CHD Risk
30-34	3%	1%	2%
35-39	5%	4%	3%
40-44	7%	4%	4%
45-49	11%	8%	4%
50-54	14%	10%	6%
55-59	16%	13%	7%
60-64	21%	20%	9%
65-69	25%	22%	11%
70-74	30%	25%	14%

* Hard CHD events exclude angina pectoris

** Low risk was calculated for a person the same age, optimal blood pressure, LDL-C 100-129 mg/dL or cholesterol 160-199 mg/dL, HDL-C ≥45 mg/dL for men or 55 mg/dL for women, non-smoker, no diabetes

Risk estimates were derived from the experience of the Framingham Heart Study, a predominantly Caucasian population in Massachusetts, USA

参考文献

- [1]. D'Agostino RB Sr, Vasan RS, Pencina MJ, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation*. 2008;117(6):743-753.